

Medication Guide

FLUOXETINE (floo ox' e teen)

CAPSULES USP

for oral use

Read the Medication Guide that comes with fluoxetine capsules before you start taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your healthcare provider about your medical condition or treatment. Talk with your healthcare provider if there is something you do not understand or want to learn more about.

What is the most important information I should know about fluoxetine?

Fluoxetine and other antidepressant medicines may cause serious side effects, including:

1. Suicidal thoughts or actions:

- **Fluoxetine and other antidepressant medicines may increase suicidal thoughts or actions** in some children, teenagers, or young adults within the **first few months of treatment or when the dose is changed**.
- Depression or other serious mental illnesses are the most important causes of suicidal thoughts or actions.
- Watch for these changes and call your healthcare provider right away if you notice:
 - New or sudden changes in mood, behavior, actions, thoughts, or feelings, especially if severe.
 - Pay particular attention to such changes when fluoxetine is started or when the dose is changed.

Keep all follow-up visits with your healthcare provider and call between visits if you are worried about symptoms.

Call your healthcare provider right away if you have any of the following symptoms, or call 911 if an emergency, especially if they are new, worse, or worry you:

- attempts to commit suicide
- acting on dangerous impulses
- acting aggressive or violent
- thoughts about suicide or dying
- new or worse depression
- new or worse anxiety or panic attacks
- feeling agitated, restless, angry or irritable
- trouble sleeping
- an increase in activity or talking more than what is normal for you
- other unusual changes in behavior or mood

Call your healthcare provider right away if you have any of the following symptoms, or call 911 if an emergency. Fluoxetine may be associated with these serious side effects:

2. Serotonin Syndrome. This condition can be life-threatening and may include:

- agitation, hallucinations, coma or other changes in mental status
- coordination problems or muscle twitching (overactive reflexes)
- racing heartbeat, high or low blood pressure
- sweating or fever
- nausea, vomiting, or diarrhea
- muscle rigidity
- dizziness
- flushing
- tremor
- seizures

3. Severe allergic reactions:

- trouble breathing
- swelling of the face, tongue, eyes or mouth
- rash, itchy welts (hives) or blisters, alone or with fever or joint pain

4. Abnormal bleeding: Fluoxetine and other antidepressant medicines may increase your risk of bleeding or bruising, especially if you take the blood thinner warfarin (Coumadin®, Jantoven®), a non-steroidal anti-inflammatory drug (NSAIDs, like ibuprofen or naproxen), or aspirin.

5. Visual problems:

- eye pain
- changes in vision
- swelling or redness in or around the eye

Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are.

6. Seizures or convulsions

7. Manic episodes:

- greatly increased energy

- severe trouble sleeping
 - racing thoughts
 - reckless behavior
 - unusually grand ideas
 - excessive happiness or irritability
 - talking more or faster than usual
8. **Changes in appetite or weight.** Children and adolescents should have height and weight monitored during treatment.
9. **Low salt (sodium) levels in the blood.** Elderly people may be at greater risk for this. Symptoms may include:
- headache
 - weakness or feeling unsteady
 - confusion, problems concentrating or thinking or memory problems
10. **Changes in the electrical activity of your heart (QT prolongation and ventricular arrhythmia including Torsades de Pointes). This condition can be life threatening. The symptoms may include:**
- fast, slow, or irregular heartbeat
 - shortness of breath
 - dizziness or fainting

Do not stop fluoxetine without first talking to your healthcare provider. Stopping fluoxetine too quickly may cause serious symptoms including:

- anxiety, irritability, high or low mood, feeling restless or changes in sleep habits
- headache, sweating, nausea, dizziness
- electric shock-like sensations, shaking, confusion

What is fluoxetine?

Fluoxetine is a prescription medicine used to treat depression. It is important to talk with your healthcare provider about the risks of treating depression and also the risks of not treating it. You should discuss all treatment choices with your healthcare provider.

Fluoxetine is used to treat:

- Major Depressive Disorder (MDD)
- Obsessive Compulsive Disorder (OCD)
- Bulimia Nervosa*
- Panic Disorder*
- Depressive episodes associated with Bipolar I Disorder, taken with olanzapine (Zyprexa)
- Treatment Resistant Depression (depression that has not gotten better with at least 2 other treatments), taken with olanzapine (Zyprexa)*

*Not approved for use in children

Talk to your healthcare provider if you do not think that your condition is getting better with fluoxetine treatment.

Who should not take fluoxetine?

Do not take fluoxetine if you:

- are allergic to fluoxetine hydrochloride or any of the ingredients in fluoxetine hydrochloride capsules. See the end of this Medication Guide for a complete list of ingredients in fluoxetine hydrochloride capsules.
- take a Monoamine Oxidase Inhibitor (MAOI). Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid.
 - Do not take an MAOI within 5 weeks of stopping fluoxetine, unless directed to do so by your physician.
 - Do not start fluoxetine if you stopped taking an MAOI in the last 2 weeks, unless directed to do so by your physician.

People who take fluoxetine close in time to an MAOI may have serious or even life-threatening side effects. Get medical help right away if you have any of these symptoms:

- high fever
- uncontrolled muscle spasms
- stiff muscles
- rapid changes in heart rate or blood pressure
- confusion
- loss of consciousness (pass out)
- **take Mellaril® (thioridazine). Do not take Mellaril® within 5 weeks of stopping fluoxetine because this can cause serious heart rhythm problems or sudden death.**
- **take the antipsychotic medicine pimozide (Orap®) because this can cause serious heart problems.**

What should I tell my healthcare provider before taking fluoxetine? Ask if you are not sure.

Before starting fluoxetine, tell your healthcare provider if you:

- Are taking certain drugs or treatments such as:
 - Triptans used to treat migraine headache
 - Medicines used to treat mood, anxiety, psychotic or thought disorders, including tricyclics, lithium, buspirone, SSRIs, SNRIs, MAOIs or antipsychotics
 - Amphetamines
 - Tramadol and fentanyl
 - Over-the-counter supplements such as tryptophan or St. John's Wort
 - Electroconvulsive therapy (ECT)

- have liver problems
- have kidney problems
- have heart problems
- have or had seizures or convulsions
- have bipolar disorder or mania
- have low sodium levels in your blood
- have a history of a stroke
- have high blood pressure
- have or had bleeding problems
- are pregnant or plan to become pregnant. Taking fluoxetine late in pregnancy may lead to an increased risk of certain problems in your newborn. Talk to your healthcare provider about the benefits and risks of treating depression during pregnancy.
 - If you become pregnant while taking fluoxetine, talk to your healthcare provider about registering with the National Pregnancy Registry for Antidepressants. You can register by calling 1-844-405-6185 or go to <https://womensmentalhealth.org/clinical-and-researchprograms/pregnancyregistry/antidepressants/>.
- are breast-feeding or plan to breast-feed. Fluoxetine may pass into your breast milk. Talk to your healthcare provider about the best way to feed your baby if taking fluoxetine.

Tell your healthcare provider about all the medicines that you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Fluoxetine and some medicines may interact with each other, may not work as well, or may cause serious side effects.

Your healthcare provider or pharmacist can tell you if it is safe to take fluoxetine with your other medicines. Do not start or stop any medicine while taking fluoxetine without talking to your healthcare provider first.

If you take fluoxetine, you should not take any other medicines that contain fluoxetine hydrochloride including:

- | |
|--|
| <ul style="list-style-type: none"> • Symbyax • Sarafem |
|--|

How should I take fluoxetine?

- Take fluoxetine exactly as prescribed. Your healthcare provider may need to change the dose of fluoxetine until it is the right dose for you.
- Fluoxetine may be taken with or without food.
- If you miss a dose of fluoxetine, take the missed dose as soon as you remember. If it is almost time for the next dose, skip the missed dose and take your next dose at the regular time. Do not take two doses of fluoxetine at the same time.
- If you take too much fluoxetine, call your healthcare provider or poison control center right away, or get emergency treatment.

What should I avoid while taking fluoxetine?

Fluoxetine can cause sleepiness or may affect your ability to make decisions, think clearly, or react quickly. You should not drive, operate heavy machinery, or do other dangerous activities until you know how fluoxetine affects you. Do not drink alcohol while using fluoxetine.

What are the possible side effects of fluoxetine?

Fluoxetine may cause serious side effects, including:

- See “What is the most important information I should know about fluoxetine?”
- **Problems with blood sugar control.** People who have diabetes and take fluoxetine may have problems with low blood sugar while taking fluoxetine. High blood sugar can happen when fluoxetine is stopped. Your healthcare provider may need to change the dose of your diabetes medicines when you start or stop taking fluoxetine.
- **Feeling anxious or trouble sleeping**

Common possible side effects in people who take fluoxetine include:

- unusual dreams
- sexual problems
- loss of appetite, diarrhea, indigestion, nausea or vomiting, weakness, or dry mouth
- flu symptoms
- feeling tired or fatigued
- change in sleep habits
- yawning
- sinus infection or sore throat
- tremor or shaking
- sweating
- feeling anxious or nervous
- hot flashes
- rash

Other side effects in children and adolescents include:

- increased thirst
- abnormal increase in muscle movement or agitation
- nose bleed
- urinating more often
- heavy menstrual periods
- possible slowed growth rate and weight change. Your child's height and weight should be monitored during treatment with fluoxetine.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of fluoxetine. For more information, ask your healthcare provider or pharmacist.

CALL YOUR DOCTOR FOR MEDICAL ADVICE ABOUT SIDE EFFECTS. YOU MAY REPORT SIDE EFFECTS TO THE FDA AT 1-800-FDA-1088.

How should I store fluoxetine?

- Store at 25°C (77°F); excursions permitted to 15° to 30°C (59 to 86°F), (See USP Controlled Room Temperature).
- Keep fluoxetine away from light.
- Keep fluoxetine bottle closed tightly.

Keep fluoxetine and all medicines out of the reach of children.

General information about fluoxetine

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use fluoxetine for a condition for which it was not prescribed. Do not give fluoxetine to other people, even if they have the same condition. It may harm them.

This Medication Guide summarizes the most important information about fluoxetine. If you would like more information, talk with your healthcare provider. You may ask your healthcare provider or pharmacist for information about fluoxetine that is written for healthcare professionals.

For more information about fluoxetine call Alembic Pharmaceuticals Limited at 1-866-210-9797.

What are the ingredients in fluoxetine?

Active ingredient: fluoxetine hydrochloride

Inactive ingredients in capsules: lactose monohydrate, microcrystalline cellulose, colloidal silicone dioxide, gelatin, titanium dioxide, FD & C Blue No. 1 and FD & C Red No. 40.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Medication Guide available at <http://www.alembicusa.com/medicationguide.aspx> or call 1-866-210-9797.

Manufactured by:

Alembic Pharmaceuticals Limited

(Formulation Division),

Panelav 389350, Gujarat, India

Manufactured for:

Alembic Pharmaceuticals, Inc.

750 Route 202, Bridgewater, NJ 08807

USA

Symbyax® (olanzapine and fluoxetine hydrochloride capsules), Sarafem® (fluoxetine hydrochloride capsules), Zyprexa® (olanzapine) are registered trademark of Eli Lilly and Company.

Sarafem® (fluoxetine hydrochloride tablets) is a registered trademark of Warner Chilcott Company, Mellaril® (thioridazine) is a registered trademark of Novartis AG Corporation, Orap® (pimozid) is a registered trademark of Teva Pharmaceuticals USA, Coumadin® (warfarin) is a registered trademark of Bristol Myers Squibb and Jantoven® (warfarin) is a registered trademark of Upsher-Smith Laboratories Inc.

Revised: 06/2020

20030359