

## Medication Guide

# Teriflunomide (ter" i floo' noe mide) Tablets, for oral use

Read this Medication Guide before you start using teriflunomide tablets and each time you get a refill. There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or your treatment.

### What is the most important information I should know about teriflunomide tablets?

#### Teriflunomide tablets may cause serious side effects, including:

- **Liver problems:** Teriflunomide tablets may cause serious liver problems, including liver failure that can be life-threatening and may require a liver transplant. Your risk of developing serious liver problems may be higher if you already have liver problems or take other medicines that also affect your liver. Your healthcare provider should do blood tests to check your liver:

- o within 6 months before you start taking teriflunomide tablets
- o 1 time a month for 6 months after you start taking teriflunomide tablets

Call your healthcare provider right away if you have any of the following symptoms of liver problems:

- o nausea
- o vomiting
- o stomach pain
- o loss of appetite
- o tiredness
- o your skin or the whites of your eyes turn yellow
- o dark urine
- **Harm to your unborn baby:** Teriflunomide tablets may cause harm to your unborn baby. Do not take teriflunomide tablets if you are pregnant. Do not take teriflunomide tablets unless you are using effective birth control.
  - o If you are a female, you should have a pregnancy test before you start taking teriflunomide tablets. Use effective birth control during your treatment with teriflunomide tablets.
  - o After stopping teriflunomide tablets, continue using effective birth control until you have blood tests to make sure your blood levels of teriflunomide tablets are low enough.
  - o If you become pregnant while taking teriflunomide tablets or within 2 years after you stop taking it, tell your healthcare provider right away.
  - o **For men taking teriflunomide tablets:**
    - If your female partner plans to become pregnant, you should stop taking teriflunomide tablets and ask your healthcare provider how to quickly lower the levels of teriflunomide tablets in your blood.
    - If your female partner does not plan to become pregnant, you and your female partner should use effective birth control during your treatment with teriflunomide tablets. Teriflunomide tablets remains in your blood after you stop taking it, so continue using effective birth control until teriflunomide tablets blood levels have been checked and they are low enough.

**Teriflunomide tablets may stay in your blood for up to 2 years after you stop taking it. Your healthcare provider can prescribe a medicine to help lower your blood levels of teriflunomide tablets more quickly. Talk to your healthcare provider if you want more information about this.**

### What is teriflunomide tablet?

- Teriflunomide tablet is a prescription medicine used to treat relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.
- It is not known if teriflunomide tablets are safe and effective in children.

### Who should not take teriflunomide tablets?

#### Do not take teriflunomide tablet if you:

- have severe liver problems.
- are pregnant or are of childbearing age and not using effective birth control.
- have had an allergic reaction to leflunomide, teriflunomide or any other ingredients in teriflunomide tablets. Please see the end of this Medication Guide for a complete list of ingredients in teriflunomide tablets.
- take a medicine called leflunomide.

### Before you take teriflunomide tablets, tell your healthcare provider about all of your medical conditions, including if you:

- have liver or kidney problems.
- have a fever or infection, or you are unable to fight infections.
- have numbness or tingling in your hands or feet that is different from your MS symptoms.
- have diabetes.
- have had serious skin problems when taking other medicines.
- have breathing problems.
- have high blood pressure.
- have had recent surgery or are planning to have surgery.
- you have an unhealed wound.

- are breastfeeding or plan to breastfeed. It is not known if teriflunomide passes into your breast milk. You and your healthcare provider should decide if you will take teriflunomide tablets or breastfeed. You should not do both.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using teriflunomide tablets and other medicines may affect each other causing serious side effects. Teriflunomide tablets may affect the way other medicines work, and other medicines may affect how teriflunomide tablets works.

Especially tell your healthcare provider if you take medicines that could raise your chance of getting infections, including medicines used to treat cancer or to control your immune system.

Ask your healthcare provider or pharmacist for a list of these medicines if you are not sure.

Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

### How should I take teriflunomide tablets?

- Take teriflunomide tablets exactly as your healthcare provider tells you to take it.
- Take teriflunomide tablets 1 time each day.
- Take teriflunomide tablets with or without food.

### What are the possible side effects of teriflunomide tablets?

**Teriflunomide tablets may cause serious side effects, including:**

- see **“What is the most important information I should know about teriflunomide tablets?”**

- **decreases in your white blood cell count.** Your white blood cell counts should be checked before you start taking teriflunomide tablets. When you have a low white blood cell count you:

- o **may have more frequent infections.** You should have a skin test for TB (tuberculosis) before you start taking teriflunomide tablets. Tell your healthcare provider if you have any of these symptoms of an infection:

- fever
- tiredness
- body aches
- chills
- nausea
- vomiting

- o **should not receive certain vaccinations during your treatment** with teriflunomide tablets and for 6 months after your treatment with teriflunomide tablets ends.

- **allergic reactions.** Stop taking teriflunomide tablets and call your healthcare provider right away or get emergency medical help if you have difficulty breathing, itching, swelling on any part of your body including in your lips, eyes, throat, or tongue.

- **serious skin reactions.** Teriflunomide tablets may cause serious skin reactions that may lead to death. Stop taking teriflunomide tablets and call your healthcare provider right away or get emergency medical help if you have any of the following symptoms: rash or redness and peeling, mouth sores or blisters.

- **other types of allergic reactions or serious problems that may affect different parts of the body such as your liver, kidneys, heart, or blood cells.** You may or may not have a rash with these types of reactions. Other symptoms you may have are:

- o severe muscle pain
- o swollen lymph glands
- o swelling of your face
- o unusual bruising or bleeding
- o weakness or tiredness
- o yellowing of your skin or the white part of your eyes

If you have a fever or rash with any of the above symptoms, stop taking teriflunomide tablets and call your healthcare provider right away.

- **ulcers of the skin or mucous membranes (such as the eyes, nose, lips, mouth, or genitals) and impaired wound healing.** Teriflunomide tablets may cause ulcers and impaired wound healing. Tell your healthcare provider right away

if any of the following happen:

- o you develop ulcers during treatment with teriflunomide tablets
- o you experience impaired wound healing during treatment with teriflunomide tablets
- o you are going to have surgery
- o you still have an unhealed wound following surgery

- **numbness or tingling in your hands or feet that is different from your MS symptoms.** You have a higher chance of getting these symptoms if you:

- o are over 60 years of age
- o take certain medicines that affect your nervous system
- o have diabetes

Tell your healthcare provider if you have numbness or tingling in your hands or feet that is different from your MS.

- **high blood pressure.** Your healthcare provider should check your blood pressure before you start taking teriflunomide tablets and while you are taking teriflunomide tablets.
- **new or worsening breathing problems.** These may be serious and lead to death. Call your healthcare provider right away or get emergency medical help if you have shortness of breath or coughing with or without fever.

The most common side effects of teriflunomide tablets include:

- headache
- diarrhea
- nausea
- hair thinning or loss (alopecia)
- increases in the results of blood tests to check your liver function

These are not all the possible side effects of teriflunomide tablets. For more information, ask your healthcare provider or pharmacist.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

#### **How should I store teriflunomide tablets?**

- Store teriflunomide tablets 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].
- Keep teriflunomide tablets and all medicines out of the reach of children.
- Bottle of 30's count comes in a child-resistant package.

#### **General information about the safe and effective use of teriflunomide tablets.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use teriflunomide tablets for a condition for which it was not prescribed. Do not give teriflunomide tablets to other people, even if they have the same symptoms you have. It may harm them.

You can ask your healthcare provider or pharmacist for information about teriflunomide tablets that is written for health professionals.

#### **What are the ingredients in teriflunomide tablets?**

**Active ingredient:** teriflunomide

**Inactive ingredients in 7 mg and 14 mg tablets:** colloidal silicon dioxide, corn starch, hydroxypropyl cellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, and sodium starch glycolate. The film coating for the 14 mg tablet is made of FD&C Blue#2 Aluminium Lake, hypromellose, polyethylene glycol 8000, talc, and titanium dioxide.

In addition, the 7 mg tablets also contain iron oxide yellow.

For more information, call Alembic Pharmaceuticals Limited at 1-866-210-9797.

All trademarks are property of their respective owners.

Manufactured by:

**Alembic Pharmaceuticals Limited**

(Formulation Division),

Panelav 389350, Gujarat, India

Manufactured for:

**Alembic Pharmaceuticals, Inc.**

Bedminster, NJ 07921, USA

20047243

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: 03/2026